



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

By following out the method indicated above we get

$$V = \frac{2}{3} R^3 [(1+e^2)E(e) - (1-e^2)F(e)].$$

COROLLARY 2. By Prof. Henry Heaton, Atlantic, Iowa.

If $R=r$, $V=1/\sin\alpha \int_0^{R-\frac{1}{2}c} \sqrt{R^2-(x-\frac{1}{2}c)^2} \sqrt{R^2-(x+\frac{1}{2}c)^2} dx$, where $x+\frac{1}{2}c$

is the distance from axis of cylinder to the plane.

Put $x=(R-\frac{1}{2}c)\sin\theta$, and $(R-\frac{1}{2}c)/(R+\frac{1}{2}c)=e$.

$$\begin{aligned} \text{Then } V &= (2/\sin\alpha)(R-\frac{1}{2}c)^2(R+\frac{1}{2}c) \int_0^{\frac{1}{2}\pi} \cos^2\theta \sqrt{1-e^2\sin^2\theta} d\theta \\ &= (2/\sin\alpha)(R+\frac{1}{2}c)^3 [(1+e^2)E(e, \frac{1}{2}\pi) - (1-e^2)F(e, \frac{1}{2}\pi)]. \end{aligned}$$

COROLLARY 3. By Prof. G. B. M. Zerr, Lebanon, Va., and Prof. C. W. M. Black, Wilbraham, Mass.

If $R=r$ and $c=0$, $V=(16r^3)/(3\sin\alpha)$.

II. Solution by G. B. M. ZERR, A. M., Ph. D., President and Professor of Mathematics in Russell College, Lebanon, Va.

Let $x^2+z^2=R^2$, be the equation to the cylinder.

Then $(x\cos\alpha+y\sin\alpha)^2+(z-c)^2=R^2$, is the equation to the auger hole.

The limits of y are,

$$y = \frac{\sqrt{R^2-(z-c)^2} - x\cos\alpha}{\sin\alpha} \quad \text{and} \quad y = -\frac{\sqrt{R^2-(z-c)^2} + x\cos\alpha}{\sin\alpha}.$$

$$\begin{aligned} \therefore V &= (4/\sin\alpha) \int_0^R \int_{-\sqrt{R^2-x^2}}^{\sqrt{R^2-x^2}} \sqrt{R^2-(z-c)^2} dz dx \\ &= (2/\sin\alpha) \int_0^R \left[\left(\sqrt{R^2-x^2} - c \right) \sqrt{R^2-(\sqrt{R^2-x^2}-c)^2} \right. \\ &\quad \left. + R^2 \sin^{-1} \left(\frac{\sqrt{R^2-x^2}-c}{R} \right) + \left(\sqrt{R^2-x^2} + c \right) \sqrt{R^2-(\sqrt{R^2-x^2}+c)^2} \right. \\ &\quad \left. + R^2 \sin^{-1} \left(\frac{\sqrt{R^2-x^2}+c}{R} \right) \right] dx. \end{aligned}$$

This does not appear to be easy to reduce.

64. Proposed by E. S. LOOMIS, A. M., Ph. D., Professor of Mathematics in Cleveland West High School, Berea, Ohio.

Find the volume and surface generated by revolving about Y , the catenary

$$y = \frac{1}{2}a(e^{x/a} + e^{-x/a}), \text{ from } x=0 \text{ to } x=a.$$

[Osborne's Calculus, page 255, example 8].

I. Solution by S. ELMER SLOCUM, Professor of Mathematics Union College, Schenectady, New York; J. M. BANDY, A. M., Professor of Mathematics in Trinity College, Trinity, N. C.; COOPER D. SCHMITT, A. M., Professor of Mathematics in the University of Tennessee, Knoxville, Tenn.; and B. F. SINE, Principal of Rock Enon High School, Rock Enon Springs, Va.

$$V = \pi \int_0^a x^2 dy. \quad y = \frac{1}{2}a(e^{x/a} + e^{-x/a}). \quad \therefore dy = \frac{1}{2}(e^{x/a} - e^{-x/a})dx.$$

$$\therefore V = \frac{1}{2}\pi \int_0^a x^2 (e^{x/a} - e^{-x/a})dx.$$

Integrating by parts,

$$V = \frac{1}{2}\pi \left[(ax^2 e^{x/a} - 2a^2 x e^{x/a} + 2a^3 e^{x/a}) + (ax^2 e^{-x/a} + 2a^2 x e^{-x/a} + 2a^3 e^{-x/a}) \right]_0^a.$$

$$\therefore V = \frac{1}{2}\pi [a^3 e + 5a^3 e^{-1} - 4a^3] = \frac{1}{2}\pi a^3 [e + 5e^{-1} - 4].$$

$$S = 2\pi \int x ds = 2\pi \left[xs - \int s dx \right],$$

where s = length of arc of catenary, $= \frac{1}{2}a(e^{x/a} - e^{-x/a})$.

$$\therefore S = 2\pi \left[\frac{1}{2}ax(e^{x/a} - e^{-x/a}) - \frac{1}{2}a \int (e^{x/a} - e^{-x/a})dx \right]_0^a$$

$$\text{or } S = 2\pi \left[\frac{1}{2}ax(e^{x/a} - e^{-x/a}) - \frac{1}{2}a^2(e^{x/a} + e^{-x/a}) \right]_0^a.$$

$$\therefore S = 2\pi [\frac{1}{2}a^2(e - 1/e) - \frac{1}{2}a^2(e + 1/e) + \frac{1}{2}2a^2]. \quad S = 2\pi a^2(1 - e^{-1}).$$

II. Solution by J. OWEN MAHONEY, B. E., M. Sc., Graduate Fellow and Assistant in Mathematics, Vanderbilt University, Nashville, Tenn.; and HENRY HEATON, M. Sc., Atlantic, Iowa.

(a). The volume is given by the equation

$$V = \pi \int_0^a x^2 dx = \frac{1}{2}\pi \int_0^a x^2 (e^{x/a} - e^{-x/a})dx.$$

Let $x = ay$, then the limits of y are 0 and 1.

$$\text{Hence } V = \pi \int_0^a x^2 dy = \frac{1}{2}\pi a^3 \int_0^1 y^2 (e^y - e^{-y})dy.$$

Integrating by parts, we find,

$$\int y^2 e^y dy = (y^2 - 2y + 2)e^y, \text{ and } \int y^2 e^{-y} dy = -(y^2 + 2y + 2)e^{-y}.$$

$$\text{Therefore } V = \frac{\pi a^3}{2} \left[(y^2 - 2y + 2)e^y + (y^2 + 2y + 2)e^{-y} \right]_0^1$$

$$= (\pi a^3 / 2)(e + 5e^{-1} - 4).$$

(b). The surface is given by

$$S = 2\pi \int_0^a x \left[1 + \left(\frac{dy}{dx} \right)^2 \right]^{\frac{1}{2}} dx = \pi \int_0^a x (e^{x/a} + e^{-x/a}) dx$$

$$= \pi a \left[e^{x/a}(x - a) - e^{-x/a}(x + a) \right]_0^a = 2\pi a^2(1 - e^{-1}).$$